MALAYSIAN NATIONAL NEONATAL REGISTRY (CRF 2011)

Supplementary Form

Instruction:

- 1) For term babies please fill in according to the most pertinent underlying cause of death.

 2) For preterm babies please fill in according to the most immediate cause of death.

| 1. Centre Name: | | Office // |
|---------------------------------|-----------|-------------------|
| 2. Name: | 3. RN: | use: Centre: |
| 4. Mother's I/C Number: New IC: | Passport: | |

| Immediate cause of death (Modified Wiggleswort | h): Tick relevant button to read | h correct classification |
|--|---|---|
| | NEONATAL DEATH | Note: LCM = Lethal Congenital Malformation |
| | (Is there any LCM? |) |
| LCM present | | © LCM absent b) (Is gestation <37 weeks?) |
| a) Lethal congenital malformation/defect, specify: Neural tube defects Anencephaly Encephalocoele Others,specify (Refer to ICD 10): CVS Complex/ cyanotic heart disease Acyanotic CNS Hydrocephalus Hydrancephaly Holoprosencephaly Others,specify (Refer to ICD 10): Recognisable syndrome Down Edward Patau Others,specify (Refer to ICD 10): | c) Gestation <37 weeks conditions associated with immaturity IVH Septicaemia PDA in failure Pulmonary hemorrhage NEC Pneumonia PIE / BPD Preumothorax Extreme prematurity Asphyxia | Gestation ≥37 weeks (Did the baby have an asphyxial condition?) d) Asphyxial condition absent (Did the baby die from infection?) e) Infection present Group B streptococcal septicaemia Meningitis Congenital pneumonia Congenital Infection Others, specify |
| Not recognisable syndrome Skeletal dysplasia Respiratory (eg. lung hypoplasia) GIT Hydrops foetalis Renal Others, specify: | | f) Other specific causes: Kernicterus/ severe neonatal jaundice Haemorrhagic disease of newborn/ Vitamin K deficiency Intracranial bleed / SAH Pneumothorax Pulmonary hemorrhage IEM MAS Surgical, specify: Others, specify: |

| Name : | Signature: | Date: | | (dd/mm/yy |
|--------|------------|-------|--|-----------|
| | | | | |